

RESTRICTED USE PESTICIDE

DUE TO ACUTE ORAL AND DERMAL TOXICITY; AND BIRD TOXICITY. FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION.

SECTION 24[C] REGISTRATION

**FOR DISTRIBUTION AND USE ONLY WITHIN
IDAHO, OREGON, AND WASHINGTON
(DO NOT USE IN LAKE COUNTY, OREGON)**



PHORATE 20 G

**EPA REG. NO. 34704-259
EPA SLN NO. ID-910014
EPA SLN NO. OR-890005
EPA SLN NO. WA-870010**

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS
ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**THIS LABEL MUST BE IN POSSESSION
OF USER AT THE TIME OF APPLICATION.**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

POTATOES — For control of aphids, leafhoppers, leaf miners, psyllids, suppression of wireworms, flea beetle larvae, and reduction of flea beetle adults. On sandy or light soils, use 11¹/₄ ozs. PHORATE 20 G per 1000 feet of row (minimum 32 inch spacing). On heavy or clay soils, use 17¹/₄ ozs. PHORATE 20 G per 1000 feet of row (any row spacing). Distribute the granules evenly in the furrow or granules may be banded on each side of the row at planting time, or at postemergence. Do not exceed 3 lbs. ai/acre per crop. Wait 90 days after treatment before harvesting potatoes.

24[C] Registrant:
Platte Chemical Co.
150 So. Main St.
Fremont, NE 68025